அற்proved For Release 1999/09/10 : CIA-RDP83-00423R001201180002-6் ு

December 4, 1953

SUBJECT: New Products - November 1953

Domestic_

Achromycin (New An	tibiotic)
Description:	Each capsule contains: Tetracycline 50 mg., 100 mg. or 250 mg.
	Each teaspoonful of Spersoids(R) dispersible powder contains: Tetracycline 50 mg. Intravenous: Tetracycline with Vitamin C 100 mg., 250 mg.
	or 500 mg.
Administration:	Recommended dosage for adults is 1 Gram per day in divided doses.
Source:	Lederle Laboratories, New York, New York.
Benetrycin Nasal S	ray (New Dosage Form)
Description:	Each pack contains: dl Desoxyephedrine 0.125%
	Pyrilamine maleate 0.15 %
	Tyrothricin 0.02 %
	Cetalkonium chloride 0.04 %
	Cotyl demethylbenzyl
Indications:	ammonium chloride
Administration:	Temporary relief of stuffed or irritated nose.
Adminis M.S. C. Coll.	Adults and children over 6, 1 or 2 sprays in each nostril every 3 to 4 hours.
Form Supplied:	Each pack contains 20 cc.
Source:	Supreme Pharmaceutical Company, New York, New York.
G	
Crysdimyein A.S. (1	
Descriptions	Each dose contains: Procaine penicillin G 600,000 u.
	Streptomycin sulfate 0.5 Gm. Dihydrostreptomycin sulfate 0.5 Gm.
	Dihydrostreptomycin sulfate 0.5 Gm. in an aqueous solution.
Indications:	Effective against gram-negative and gram-positive
	organisms; valuable in mixed or undiagnosed infections,
Administration:	Maximum: 3 cc. Minimum: 13 cc.
Form Supplied:	Vials of 5 dose of 3 cc. each.
Source:	E.R. Squibb and Sons, New York, New York.

Deltamide W/Penicillin (New Form)

Description: Fach tablet or teaspoonful contains:

Penicillin 250,000 u.
Sulfadiazine 0.167 Gm.
Sulfamerazine 0.167 Gm.
Sulfamethazine 0.056 Gm.
Sulfacetamide 0.111 Gm.

Tablets are buffered with calcium carbonate and the chocolate-flavored suspension with sodium citrate.

Source: Armour Laboratories, Chicago, Illinois.

*Dihydrostreptomycin-Streptomycin (Parenteral) (New Dosage Form)
Description: Each vial contains: Streptomycin sulfate

Dihydrostreptomycin sulfate

in equal quantities.

Form Supplied: 1 Gram vials and 5 Gram vials.

Source: Parke, Davis and Company, Detroit, Michigan.

**Dipen (New Product)

Description: Contains: Penicillin proceins 300,000 u.

Potassium panicillin buffered 100,000 u.
Dihydrostreptomycin sulfate 0.5 Gram or

1.0 Gram

Form Supplied: Packaged in 1 dose and 5 dose gizes.

Source: Standard Pharmacoutical Company, Inc., New York, New York.

K-Cillin (New Product)

Description: Each 5 cc. contains: Penicillin 300,000 or 600,000 u.

in stable suspension.

Said to possess a 3 year "shelf life".

Administration: Orally.

Form Supplied: 2 oz. bottles.

Source: Bic-Ramo Drug Company, Baltimore, Maryland.

Pon-G-Cap (New Dosage Form)

Description: Each capsule contains: Penicillin G 200,000 u.

suspended in oil.

Indications: To maintain prolonged blood levels. Form Supplied: Bottles of 12, 100 and 500 capsules.

Source: The Upjohn Company, Kalamazoo, Michigan.

	mo	

(New Product) *Bi-Gor-Mone Description: Contains: Testosterone propionate 25 mg. 1 mg. Estradiol benzoate in sesame oil. Indications: Menopause, functional uterine bleeding, inhibition of lactation and in geriatrics. 10 cc. vials. Form Supplied: C.F. Kirk, New York, New York. Source: Gynetone Repetabe (New Form-Repetab) Description: Each ".02" repetab contains: Ethinyl estradiol. 0.02 mg. Methyl testosterone 5.0 mg. Each ".04" repetab contains: Ethinyl estradiol O.L mg. Methyl testosterone 10.0 mg. For treatment of the menopause and osteoporosis Indications: as well as for the management of geriatric patients where hormonal supplementation is desirable. Monopause: 1 or 2 ".04" or 2 to 4 ".02" repetabs daily. Administration: Maintenance dosage 1 ".02" repetat daily. Osteoporosis and in geriatrics: 1 or 2 ".02" daily. Bottles of 30 and 100 repatabs. Form Supplied: Source: Schering Corporation, Bloomfield, New Jersey. Thecandrol Oral. (New Dosage Form) Description: Each capsule contains: Ketohydroxyestratiene (Menagen) 10,000 i.u. Methyl testosterone 10.0 mg. Also available in half strength capsules. Indications: Symptomatic relief for menopausal patients. Average: High initial desage of 2 capsules daily for Administration: one week, followed by a maintenance dosage of 1 capsule daily.

Bottles of 100 capsules. Form Supplied: Source:

Parke, Davis and Company, Datroit, Michigan,

<u>Vitamins</u>

Pro-Mons (New	Product)						
Description:	Each	enteric	tablet	contains:	Ammonium chloride	5.0	grn.
_					Homatropine mbr.	0.5	mg.
					Caffeine	0.5	mg.
					Thiamine hol.	2.0	mg o
					Riboflavîn	1.0	_
					Pyridozine hel.	0.5	_
					Calcium pantothenate	1.0	_
					Niacinamide	5.0	_

Indications: Nervous and emotional instability, irritability, crying

spells, headaches, nausea and other symptoms of the

pre-menstrual syndrome.

Administration: 2 tablets, t.i.d., after meals, beginning 10 days

before menstrual period.

Form Supplied: Bottles of 60 tablets. School For Releases 1929 (1941); SIA-RDF, 82-00423R001201180002-6

Revicapa (New Product) Description: Each capsule contains: d-amphetamine sulfate 5 mg. Methyl cellulose 200 mg. Vitamins A, Bl. B20 B60 B120 C. D. and minerals. Indications: For the control of obesity. Administration: Average: 1 to 2 capsules 3 times daily, ½ to an hour before meals. Form Supplied: Bottles of 100 and 1,000 capsules. Source: Lederle Laboratories, New York, New York, Thienicavin (New Product) Description: High potency form of injectable vitamin B complex. In beriberi, pellagra, anorexia, impaired lactation Indications: and pernicious vemiting of pregnancy, arthritis, gout, polyneuritis, functional gastro-intestinal disturbances, hyperthyroidism, tachycardia, palpitation and avitaminosis B. Form Supplied: Vials of 30 cc. Chicago Pharmacal, Chicago, Illinois. Source: T.V.D. Formula (New Product) Description: Each tablet contains: Thyroid USP 10 mg. Thiamine hel. 2 mg. Riboflavin 2 mg. Niacinamide 10 mga Ascorbic acid 20 mg. d-amphetamine phosphate 5 mg. Indications: As an adjunct to dietary treatment in the medical management of obssity to lessen the desire for food; to supply more than the minimum daily requirements of Vitamins B and C and to replace any loss in dieting; to provide calorgenic action and to help control water balance. 1 tablet one hour before each meal, Administration: Form Supplied: Bottles of 50 and 160 tablets. Sources Adams Laboratories, Buffalo, New York, <u>Antihistamines</u> Benacine (New Product) Description: Each tablet contains: Benadryl 25₀0 Hyosoine hydrobromide 0.325 mg. Indications: The therapy of motion sickness and of parkinsonism. Form Supplied: Bottles of 100 and boxes of 12.

Parks, Davis and Company, Ltd., Detroit, Michigan.

Source

Approved For Release 1999/09/10 : CIA-RDP83-00423R001201180002-6

-5a

Triohistin (New Product)

Description: Each triple tablet contains:

Methapyrilene hydrochloride

10 mg. 10 mg.

Pyrilamine maleate

TO mão

Phenyltoloxamine dihydrogen clirate

10 mg.

Indications:

Relief of the discomforts of hay fever, vasomotor

rhinitis, eczema and other allergies.

Administration:

1 tablet three times a day.

Form Supplied:

Bottles of 100 and 1,000 tablets.

Source:

B.F. Ascher and Company, Kansas City, Missouri.

Veterinary

Wyoth Laboratories recently advertised a new antibiotic treatment for chicken air-sac infections, 'Bicillin-SM', a combination of Bicillin and dihydrostreptomycin.

Correction

'Cilloral' Suspension is produced by Bristol Laboratories. The source was committed in last months' report.

The product called "Erythrocin tablets" in last months' report should be Erythromyein tablets. This product is made by The Upjohn Company while Erythrocin is the brand name for Abbott's erythromyein products.

Foreign

Hostamyein and Hostamyein Forte (New Products) Description: Hostamycin contains: Novocaine-penicillin G 400,000 u. Penicillin G sodium 100,000 u. Dihydrostreptomycin sulfate 0.5 Gm. Hostamycin Forte contains: Novocaine-penicillin G 300,000 u, Penicillin G sodium 100,000 u. Dihydrostreptomycin sulfate 0.5 Gm. In endocarditis, cholecystitis, peritonitis, Indications: puerperal sepsis, etc. Administration: Usually 1 daily injection intramuscularly is sufficient in slight infections. For more severe infections, larger quantities, divided into two or four doses per 24 hours.

Form Supplied:

1 Bottle with ampule of solvent. Hospital packings of 25 bottles.

Source:

Farbwerke Hoechst, A.G., Frankfurt-am-Main, Germany,

Intrathecal Streptomycin

Description:

Each ampule contains: Streptomycin Solution of suitable concentration is obtained

by adding 5 mils of solvent to 1 ampule.

Form Supplied:

Source:

The Distillers Company, England.

Intrathecal Streptomycin

Description:

Each ampule contains: Streptomycin 50 mgm. or 100 mgm.

Form Supplied:

Boxes of 5 ampules.

Boxes of 10 ampules.

Source:

Glaxo Laboratories, Ltd., Greenford, England.

Pendiamezine Cobaltosa (New Product)

Descriptions Each tablet contains: Potassium penicillin 150,000 u.

Cobalt 0.003 Gm. Sulfadiazino 0,167 Gm. Sulfamerazine 0.167 Gm. Sulfamotazina 0.167 Gm.

Indications:

In the treatment of and prophylaxis of infections caused by gram-positive and gram-negative germs.

Administration:

Adults: 4 to 8 tablets daily. Children: 2 to 6 tablets daily.

Form Supplied:

l aluminum wrapped tablet.

Source:

Laboratorios Life, Bogota, Colombia.

(New in Market)

Wyeth Laboratories, London, England is now marketing a Penicillin inhalation set for nasal and oral inhalation. Each set contains an inhalator and two vials, each containing three capsules of 100,000 units of crystalline penicillin potassium salt in each. The set is also supplied separately: 1 inhalator, 1 box of 6 capsules or 1 box of 9 capsules.

(New Names)

Allen and Hanburys, Ltd., London, England, is now distributing their product, Guanillin (New Products Report, June, 1953), under the new name, Guanimycin.

Bristol Laboratories is now distributing, in Nicaragua, their product, Panbiotic, under the new name of Tribiogyl. This product is competitive to Pfizer's Pronapen Plus.

Sharp and Dohme is distributing their U.S. product, Remanden, under the name of Penbenamid in the United Kingdom.

(New Strength Only)

E.R. Squibb and Sons is now distributing the U.S. products, "Cristifor \$800" and \$1200" in Nicaragua under the name of "Despacilina plus \$800" and \$1200".

Hormones

Cibagthen (New Product)

Description: Dry preparation contains: ACTH in pure state

Gel preparation contains: Cibacthen with gelatin in

a sterile solution.

Form Supplied: Dry: Bottles of 10 and 40 units.

Gel: Bottles of 2 cc. containing 40 units.

Source: Ciba, A.G., Wehr-am-Baden, Germany.

G.T. 50B (New Form)

Description: Each 5 mil ampule contains: Calciferol 13.6 mgm.

Neostigmine bromide 0.295 mgm.
Carbachol 0.1 mgm.
Ethinylestradiol 0.01 mgm.
Progesterone 5.0 mgm.
Procaine 10.0 mgm.

hydrochloride

Indications: Treatment of rheumatoid arthritis, arthrosis,

oste oporosis and oste omalcia.

Form Supplied: Packages of 5 and 20 ampules.

Source: Bengue and Company, Ltd., Wembley, England.

Perenquinol - A (New Product)

Description: Each cc. contains: Estradiol bensoate

2.5 mg, Progesterone 12.5 mg.

Indications: Administrations:

Diagnostic test of pregnancy and secondary amenorrhea. 1 ampule intramuscularly every day for two days.

Form Supplied:

Boxes of 2 ampules of 1 cc. each.

Source:

Parmox Laboratories, (Marxuach and Zegri), San Juan, P.R.

Perandren "M" Crystules (New Dosage Form)

Description: Source:

Each co. contains: Testosterone isobutyrate 50 mgm.

Ciba Laboratories, Horsham, England.

Pro-Estrin (New Product)

Descriptions

Each cc. contains: Estradiol benzoate

2.5. mg.

Indications: Administration:

Progesterone 12.5 mg. Diagnostic test of pregnancy and secondary amenorrhes. l mpules intramuscularly every day for three days.

Form Supplied: Sources

Boxes of 3 ampules of 1 cc. each. Afleg Laboratories, San Juan, P.R.

Scheroson (New Product) Description:

Each 2 cc. contains: Cortisons acetate

10 mg.

Indications:

Chloramphanicol in oil solution 4 mg. Inflammation and allergic conditions of the external

Administration:

Local Donthalmic application.

Form Supplied:

Tube of 2 co.

Source

Schoring, A.G., Berlin-West, Germany,

<u>Vitamina</u>

Appatinic Tablets (New Product)

Description:

Each sugar-coated tablet contains:

Vitamin B₁₂ (from streptomyces fermentation) Folic acid Ascorbic acid

0.4 mg. 50.0 mg.

5.0 meg.

Ferrous sulfate USP orystals Liver-stomach concentrate

194.0 mg. 100,0 mg.

Indications:

A comprehensive hematinic for those cases of anemia which do not respond to conventional iron therapy.

Administration:

2 tablets 3 times daily at mealtime. For children,

1 tablet three times daily.

Form Supplied:

Bottles of 100, 500 and 1,000 tablets.

Source:

Anglo-Canadian Drug Company, Ltd., Oshawa, Canada.

Ferrogen Compound	(New Product)	
Description:	Each chocolate-coated tablet contains:	
	Ferrous sulfate 325.0 mg.	
	Bone flour (edible) 325.0 mg.	
	Vitamin A 1,500 i.u.	
	Vitamin D 500 i.u.	
	Vitamin B, l.O mg.	
	Riboflavin 1.0 mg.	
	Nieginamido 5.0 mg.	
	Ascorbic acid (Vitamin C) 30.0 mg.	
	Sodium iodido 0,2 mg.	
Indications:	A distary supplement in pregnancy, lactation,	
	convalescence and iron deficiency anemia where	
	vitamin therapy is indicated.	
Administration:	1 tablet three times daily.	
Form Supplied:	Bottles of 100 and 500.	
Sources	Charles E. Frosst and Company, Montreal, Canada,	
	outside of the second of the second of the second of	•
Maturex Injectable	(dried) (New Product)	
Description:	Each vial contains: Vitamin Bl2 crystalline	30 mcg.
-	Thiamine	10 mg.
	Riboflavin	2 mg.
	Niacinamide	1.00 mg.
	Pyridoxine	5 mg.
	d-Panthenol	
	Vitamin B factors as derive	5 mg.
	15 Grams of fresh liver.	d Iron
Indications:	Pernicious anemia, other macrocytic anemias, vit	
	B deficiencies.	amin
Administration:	Macrocytic anemias: 1 to 2 cc. once or twice we	~1eT
	for therapeutic effect; 1 cc. daily or every sec	ekly
	day for 3 months if combined degeneration presen	ond •
Form Supplied:	Boxes of 5 x 1 dose vials with 10 cc. sterile di	. Vo 3
Source:	Ayerst, McKenna and Harrison, Ltd., Montreal, Ca	Trigue ^o
	ajorso, randina and narrison, node, rontreal, ca	nacao
Neo-Hepatrex (New	Product)	
Description:	Vitamin B12 in natural complex forms	
	Liver extract obtained by fermentative decomposi	+4 ms
Indications:	In all forms of anemia due to lack of the anti-	OTOHO
	anomic factors in the liver.	
Form Supplied:	Packs of 2 cc. ampules.	
Source:		C
would be	C.F. Boehringer and Soehne, G.m.b.H., Mannheim,	cormany.

Approved For Release 1999/09/10: CIA-RDP83-00423R001201180002-6

-10-

Ostogen-A with C (New Product) 8,000 i.u. Description: Each co. contains: Vitamin A 4,000 lou Vitamin D Vitamin C 1.20 mg. Drop-dose vitamin supplement designed especially for Indications: infants and children. Administration: 5 drops daily, mixed with milk or fruit juices or dropped directly on the tongue. Dropper bottles of 8 cc., 15cc. and 30 cc. Form Supplied: Charles Trosst and Company, Montreal, Canada. Source:

Mead Johnson and Company is now distributing their vitamin-mineral product for pregnancy stresses, <u>Natalins</u>, under the name of <u>Obelina</u> in Canada.

Veterinary

Nec-Tac Cintment N.F. (New Product)

Description: Each tube contains: Necmycin base 30 mg.

Dihydrostreptomycin base 250 mg.

Bacitracin 2,000 i.u.

Procaine penicillin G 250,000 i.u.

Phthalylsulfacetamide 10 mg.

in free-flowing monostearate base.

Indications: Treatment of bovine mastitis.

Form Supplied: Tubes containing 7.5 cc.

Source: Austin Laboratories, Ltd., Winnipeg, Canada.

KESTKTOTES	10001101	<u> </u>	DP83-004	23R0012 0118
•				
1	ROUTING AND C	CONTROL R	ECORD	
DO NOT	DETACH FROM	'ON LOAN	' DOCUME!	NTS
			-25X1A2g	:
το:	TA I hanny	· ···········		
ATTN:	IA Library			
BUILDING	·····		ROOM NO.	
TITLE				
	of new pharmad			
REMARKS				
EXXX RETAIN		ON L	DAN	
DOCUMENT(S)	DOCUMENT(S) MUST B	DOCUMENT(S) MUST BE RETURNED TO CONTACT DIVISION/OO		
FOR RETENSION BY ADDRESSEE	BY (DEADLINE)	OUTTRO	25X1/	:
FROM:	ACT DIVISION/O	BRANCH O Scient		<u> </u>
BUILDING		ROOM NO.		